



Lesson 7: Informal Proofs of Properties of Dilations

Student Outcomes

- Students know an informal proof of why dilations are angle-preserving transformations.
- Students know an informal proof of why dilations map segments to segments, lines to lines, and rays to rays.

Lesson Notes

These properties were first introduced in Lesson 2. In this lesson, students think about the mathematics behind why those statements are true in terms of an informal proof developed through a discussion. This lesson is optional.

Classwork

Discussion (15 minutes)

Begin by asking students to brainstorm what they already know about dilations. Accept any reasonable responses. Responses should include the basic properties of dilations; for example, lines map to lines, segments to segments, and rays to rays. Students should also mention that dilations are angle-preserving. Let students know that in this lesson they informally prove why the properties are true.


- In previous lessons, we learned that dilations are angle-preserving transformations. Now we want to develop an informal proof as to why the theorem is true:

THEOREM: *Dilations preserve the measures of angles.*


- We know that dilations map angles to angles. Let there be a dilation from center O and scale factor r . Given $\angle PQR$, we want to show that if $P' = \text{Dilation}(P)$, $Q' = \text{Dilation}(Q)$, and $R' = \text{Dilation}(R)$, then $|\angle PQR| = |\angle P'Q'R'|$. In other words, when we dilate an angle, the measure of the angle remains unchanged. Take a moment to draw a picture of the situation. (Give students a couple of minutes to prepare their drawings. Instruct them to draw an angle on the coordinate plane and to use the multiplicative property of coordinates learned in the previous lesson.)
 - *(Have students share their drawings.) A sample drawing is below.*

Scaffolding:

Provide more explicit directions, such as: “Draw an angle PQR , and dilate it from a center O to create an image, angle $P'Q'R'$.”



- Could the line containing $\overrightarrow{Q'P'}$ be parallel to the line containing \overrightarrow{QP} ?
 - Yes Based on what we know about the fundamental theorem of similarity, since $P' = \text{Dilation}(P)$ and $Q' = \text{Dilation}(Q)$, then we know that the line containing $\overrightarrow{Q'P'}$ is parallel to the line containing \overrightarrow{QP} .
- Could $\overrightarrow{Q'P'}$ intersect \overrightarrow{QR} ?
 - Yes, it intersects \overrightarrow{QR} (as shown in sample student drawing) or will intersect if ray $\overrightarrow{Q'P'}$ is extended (as needed in actual student work).
- Could $\overrightarrow{Q'P'}$ be parallel to \overrightarrow{QR} ?
 - No Based on what we know about the fundamental theorem of similarity, \overrightarrow{QR} and $\overrightarrow{Q'R'}$ are supposed to be parallel. In the last module, we learned that there is only one line that is parallel to a given line going through a specific point. Since $\overrightarrow{Q'P'}$ and $\overrightarrow{Q'R'}$ have a common point, Q' , only one of those lines can be parallel to \overrightarrow{QR} .
- Now that we are sure that $\overrightarrow{Q'P'}$ intersects \overrightarrow{QR} , mark that point of intersection on your drawing (extend rays if necessary). Let's call that point of intersection point B .
 - A sample student drawing is below:




- At this point, we have all the information that we need to show that $|\angle PQR| = |\angle P'Q'R'|$. (Give students several minutes in small groups to discuss possible proofs for why $|\angle PQR| = |\angle P'Q'R'|$.)

 - We know that when parallel lines are cut by a transversal, then their alternate interior angles are equal in measure. Looking first at the parallel lines containing rays $Q'P'$ and QP , we have transversal \overline{QB} . Then, alternate interior angles are equal (i.e., $|\angle Q'BQ| = |\angle PQR|$). Now, looking at the parallel lines containing rays $Q'R'$ and QR , we have transversal $\overline{Q'B}$. Then, alternate interior angles are equal (i.e., $|\angle P'Q'R'| = |\angle Q'BQ|$). We have the two equalities, $|\angle P'Q'R'| = |\angle Q'BQ|$ and $|\angle Q'BQ| = |\angle PQR|$, where within each equality is $\angle Q'BQ$. Therefore, $|\angle PQR| = |\angle P'Q'R'|$.

Scaffolding:

Remind students what they know about angles that have a relationship to parallel lines. They may need to review their work from Topic C of Module 2. Also, students may use protractors to measure the angles as an alternative way of verifying the results.

A sample drawing is below:



- Using FTS and our knowledge of angles formed by parallel lines cut by a transversal, we have proven that dilations are angle-preserving transformations.


Exercise (5 minutes)

Following this demonstration, give students the option of either (1) summarizing what they learned from the demonstration or (2) writing a proof as shown in the exercise.

Exercise

Use the diagram below to prove the theorem: *Dilations preserve the measures of angles.*

Let there be a dilation from center O with scale factor r . Given $\angle PQR$, show that since $P' = \text{Dilation}(P)$, $Q' = \text{Dilation}(Q)$, and $R' = \text{Dilation}(R)$, then $|\angle PQR| = |\angle P'Q'R'|$. That is, show that the image of the angle after a dilation has the same measure, in degrees, as the original.




Using FTS, we know that the line containing $\overline{Q'P'}$ is parallel to the line containing \overline{QP} and that the line containing $\overline{Q'R'}$ is parallel to the line containing \overline{QR} . We also know that there exists just one line through a given point, parallel to a given line. Therefore, we know that $\overline{Q'P'}$ must intersect \overline{QR} at a point. We know this because there is already a line that goes through point Q' that is parallel to \overline{QR} , and that line is $\overline{Q'R'}$. Since $\overline{Q'P'}$ cannot be parallel to $\overline{Q'R'}$, it must intersect it. We let the intersection of $\overline{Q'P'}$ and \overline{QR} be named point B . Alternate interior angles of parallel lines cut by a transversal are equal in measure. Parallel lines QR and $Q'R'$ are cut by transversal $\overline{Q'B}$. Therefore, the alternate interior angles $P'Q'R'$ and $Q'BQ$ are equal in measure. Parallel lines $\overline{Q'P'}$ and \overline{QP} are cut by transversal \overline{QB} . Therefore, the alternate interior angles PQR and $Q'BQ$ are equal in measure. Since the measures of $\angle P'Q'R'$ and $\angle PQR$ are equal to the measure of $\angle QB'Q$, then the measure of $\angle PQR$ is equal to the measure of $\angle P'Q'R'$.

Example 1 (5 minutes)

In this example, students verify that dilations map lines to lines.

- On the coordinate plane, mark two points: A and B . Connect the points to make a line; make sure you go beyond the actual points to show that it is a line and not just a segment. Now, use what you know about the multiplicative property of dilation on coordinates to dilate the points from center O by some scale factor. Label the images of the points. What do you have when you draw a line through points A' and B' ?

Have several students share their work with the class. Make sure each student explains that the dilation of line AB is the line $A'B'$. Sample student work is shown below.




- Each of us selected different points and different scale factors. Therefore, we have informally shown that dilations map lines to lines.

Example 2 (5 minutes)

In this example, students verify that dilations map segments to segments.

- On the coordinate plane, mark two points: A and B . Connect the points to make a segment. This time, make sure you do not go beyond the marked points. Now, use what you know about the multiplicative property of dilation on coordinates to dilate the points from center O by some scale factor. Label the images of the points. What do you have when you connect point A' to point B' ?




Have several students share their work with the class. Make sure each student explains that the dilation of segment AB is the segment $A'B'$. Sample student work is shown below.

- Each of us selected different points and different scale factors. Therefore, we have informally shown that dilations map segments to segments.

Example 3 (5 minutes)

In this example, students verify that dilations map rays to rays.

- On the coordinate plane, mark two points: A and B . Connect the points to make a ray; make sure you go beyond point B to show that it is a ray. Now, use what you know about the multiplicative property of dilation on coordinates to dilate the points from center O by some scale factor. Label the images of the points. What do you have when you connect A' through point B' ?



Have several students share their work with the class. Make sure each student explains that the dilation of ray AB is the ray $A'B'$. Sample student work is shown below.

- Each of us selected different points and different scale factors. Therefore, we have informally shown that dilations map rays to rays.

Closing (5 minutes)

Summarize, or ask students to summarize, the main points from the lesson.

- We know an informal proof for dilations being angle-preserving transformations that uses the definition of dilation, the fundamental theorem of similarity, and the fact that there can only be one line through a point that is parallel to a given line.
- We informally verified that dilations of segments map to segments, dilations of lines map to lines, and dilations of rays map to rays.

Exit Ticket (5 minutes)




Name _____

Date _____

Lesson 7: Informal Proofs of Properties of Dilations

Exit Ticket


Dilate $\angle ABC$ with center O and scale factor $r = 2$. Label the dilated angle, $\angle A'B'C'$.



1. If $\angle ABC = 72^\circ$, then what is the measure of $\angle A'B'C'$?
2. If the length of segment AB is 2 cm, what is the length of segment $A'B'$?
3. Which segments, if any, are parallel?

Exit Ticket Sample Solutions

Dilate $\angle ABC$ with center O and scale factor $r = 2$. Label the dilated angle, $\angle A'B'C'$.



1. If $\angle ABC = 72^\circ$, then what is the measure of $\angle A'B'C'$?

Since dilations preserve angles, then $\angle A'B'C' = 72^\circ$.

2. If the length of segment AB is 2 cm, what is the length of segment $A'B'$?

The length of segment $A'B'$ is 4 cm.

3. Which segments, if any, are parallel?


Since dilations map segments to parallel segments, then $\overline{AB} \parallel \overline{A'B'}$, and $\overline{BC} \parallel \overline{B'C'}$.

Problem Set Sample Solutions

1. A dilation from center O by scale factor r of a line maps to what? Verify your claim on the coordinate plane.

The dilation of a line maps to a line.


Sample student work is shown below.



2. A dilation from center O by scale factor r of a segment maps to what? Verify your claim on the coordinate plane.

The dilation of a segment maps to a segment.


Sample student work is shown below.



3. A dilation from center O by scale factor r of a ray maps to what? Verify your claim on the coordinate plane.

The dilation of a ray maps to a ray.

Sample student work is shown below.




4. **Challenge Problem:**

Prove the theorem: *A dilation maps lines to lines.*

Let there be a dilation from center O with scale factor r so that $P' = \text{Dilation}(P)$ and $Q' = \text{Dilation}(Q)$. Show that line PQ maps to line $P'Q'$ (i.e., that dilations map lines to lines). Draw a diagram, and then write your informal proof of the theorem. (Hint: This proof is a lot like the proof for segments. This time, let U be a point on line PQ that is not between points P and Q .)

Sample student drawing and response are below:



Let U be a point on line PQ . By the definition of dilation, we also know that $U' = \text{Dilation}(U)$. We need to show that U' is a point on line $P'Q'$. If we can, then we have proven that a dilation maps lines to lines.

By the definition of dilation and FTS, we know that $\frac{|OP'|}{|OP|} = \frac{|OQ'|}{|OQ|}$ and that \overline{PQ} is parallel to $\overline{P'Q'}$. Similarly, we

know that $\frac{|OQ'|}{|OQ|} = \frac{|OU'|}{|OU|} = r$ and that line QU is parallel to line $Q'U'$. Since U is a point on line PQ , then we also

know that line PQ is parallel to line $Q'U'$. But we already know that \overline{PQ} is parallel to $\overline{P'Q'}$. Since there can only be one line that passes through Q' that is parallel to line PQ , then line $P'Q'$ and line $Q'U'$ must coincide. That places the dilation of point U, U' , on the line $P'Q'$, which proves that dilations map lines to lines.