



Lesson 3: Analyzing a Verbal Description


Student Outcomes

- Students make sense of a contextual situation that can be modeled with a linear, quadratic, or exponential function when presented as a word problem. They analyze a verbal description and create a model using an equation, graph, or table.

Lesson Notes

This lesson asks students to recognize a function type from a verbal description of a context, using linear, quadratic, and exponential functions and linear inequalities. They formulate a model that can be used to analyze the function in its context. For this lesson, they do not go beyond the second step in the modeling cycle but focus on recognition and formulation only. Unlike in previous modules, no curriculum clues are provided (e.g., lesson or module title) to guide students toward the type of function represented by the situation. There is a mix of function types, and students learn to recognize the clues that are in the description itself. They analyze the relationship between the variables and/or the contextual situation to identify the function type.

Throughout this module, refer to the modeling cycle below (found on page 61 of the CCLS and page 72 of the CCSSM):



MP.1
&
MP.4

Throughout this lesson, students use the first steps in the modeling cycle. When presented with a problem, they make sense of the given information, define the variables involved, look for entry points to a solution, and create an equation to be used as a model for the context.

Classwork

Opening (15 minutes)

There are a number of everyday problems that are used for mathematical modeling. It may be helpful to review them quickly before beginning this lesson. Present these on the board or screen, and invite students to take notes in their math journals or notebooks.

This Opening is meant to offer suggestions for deciphering descriptions of contextual situations and is not intended to be comprehensive or a list of keywords. Students should use this information to find an entry point to solving a problem and should not be encouraged to do keyword reading of modeling problems.

Scaffolding:

Consider skipping this Opening if students do not need this review.

Rate Problems—Linear

Rate problems relate two quantities, usually in different units. Rate problems commonly use some measure of distance and time such as feet per second. These are called *uniform rate* word problems. Rate problems might also use quantities that are not related to distance or time (e.g., pills per bottle, dollars per pound). Rate problems may also relate the number of people to a space (e.g., students per class), the number of objects to a space (e.g., corn stalks per row or rows of corn per field), dollars per mile (e.g., taxi fare), or dollars per person. The possibilities are limitless, and remember that sometimes the uniform rate problem includes a flat fee plus a rate per unit, making the linear model a bit more interesting.

- The familiar formula (Rate)(Time) = Distance can be adjusted to accommodate any (or all) of the uniform rate problem situations mentioned above. These use the equation of the form $f(x) = mx$ (or $d = rt$).
- Sometimes the rate is combined with a constant part (e.g., flat fee). For example, a taxi service charges \$4 plus \$2.50 per mile (times the mileage). These equations use the form $f(x) = mx + b$.
- These problems might include inequalities with a maximum or minimum amount possible set by real-life conditions. For example, in a rate problem there may be a minimum number of corn stalks needed in a stalks-per-row problem or a maximum number of classrooms in a students-per-classroom problem.

Note: Be careful not to confuse a simple rate problem with one that involves the motion of a free falling object.

Objects in Motion and Area—Quadratic

Motion problems are related to free falling objects¹ or projectiles under the influence of gravity. The functions used to model these functions are always quadratic and relate the distance from the initial position (i.e., the height from which the object was dropped or projected into the air), measured in either feet or meters, to the time that has passed (i.e., the number of seconds since the object was dropped or projected into the air).

In area problems with dimensions that are variable and linear, expressions representing areas are quadratic.

- When distances are measured in feet, we use $h(t) = -16t^2 + v_0t + h_0$, where h represents height of the object in feet, t represents time in seconds, v_0 represents initial velocity in feet per second, and h_0 represents initial position (starting height) in feet.
- When distances are measured in meters, we use $h(t) = -4.9t^2 + v_0t + h_0$, with all the same variables as above.
- When linear measurements have first-degree variables and are used to find area, the result is likely a quadratic model.
- For rectangular area, $A = (l)(w)$; for triangles, $A = \frac{1}{2}(b)(h)$; for trapezoids, $A = \frac{1}{2}(b_1 + b_2)(h)$. All of these, and many others, can lead to a quadratic function that models the area of the 2-dimensional figure.

Growth Problems—Exponential

Exponential problems could involve growth or decay of money (e.g., interest earned or paid), percent growth or decay (e.g., inflation or depreciation), radioactive material (e.g., half-life problems), population (e.g., bacteria, humans, or rabbits), etc.

¹Free falling objects: Objects that are projected, launched, thrown, or dropped, without the aid of a motor (or other device) that provides additional force beyond the initial projection of the object. The object might be shot from a cannon or dropped from a cliff but will have no rotors, motor, wing, or other device to keep it aloft or defy gravity.

- The general form for any problem related to exponential growth is $f(x) = ab^x$, where a and b are constant values. You can use this form for just about every exponential growth problem, but there are some special cases for which we have special forms of this same function (see the following).

Except in problems of compounded interest, encourage students to use the general exponential form for most problems related to growth, as it is usually the most efficient.

- Depending on how the interest is compounded, we would use different forms of the growth function. Until you get a little further in your math studies, we will use the annual compound interest formula:

$$P(t) = P_0 \left(1 + \frac{r}{n}\right)^{nt}$$
 where P represents the current or future value of the money, P_0 is the principal amount (the initial investment), r is the annual interest rate, n represents the number of times per year the interest is compounded, and t is the number of years the interest is earned or paid.
- Population growth is generally modeled with the general exponential function form. Sometimes it is helpful to know a few commonly used variables: $P(n) = P_0(1 + r)^n$, where P represents the future population, P_0 is the initial population, r is the growth rate, and n is the number of time periods of growth.
- Sometimes in growth problems, you need to determine a percentage rate of growth over an interval of time. If that happens, use this strategy: Find the total growth for the interval, $(P - P_0)$ and divide by P_0 (the starting population or value). This gives you a ratio representing the new growth to the starting population or value. This result will be in decimal form and can be used that way or changed to a percentage. So, to recap: A percentage rate of growth for an interval can be determined by the expression: $\frac{100(P - P_0)}{P_0}$.
- Do you see that all these exponential formulas are different forms of the general exponential equation in the first bullet? $f(x) = ab^x$?

To highlight MP.7, ask students to identify and interpret a and b in the various examples above.

Have students work the three examples below with a partner or small groups. Give them about 2 minutes to work on the function, and then pause to have the group discuss the suggested questions together. Try to keep this to about 5 minutes per example.

Example 1 (5 minutes)

Have students read the problem. Then, use the questions that follow to guide a discussion. Offering a variety of tools for students to use while engaging with these examples promotes MP.5 and allows multiple entry points for model development. Consider offering access to a graphing calculator or graphing computer software, graph paper, and a spreadsheet program.

Example 1

Gregory plans to purchase a video game player. He has \$500 in his savings account and plans to save \$20 per week from his allowance until he has enough money to buy the player. He needs to figure out how long it will take. What type of function should he use to model this problem? Justify your answer mathematically.

Scaffolding:

- Have struggling students keep notes in their math journals or notebooks where all contextual scenarios (and the corresponding functions used) are recorded as the class works through the examples.
- Having this notebook readily available benefits all students.

- Gregory decides that the exponential function can best represent the situation. Do you agree or disagree? Why? Support your answer mathematically.
 - *I disagree because Gregory's money increases at a constant rate of \$20 a week with a starting balance of \$500. The graph of the amount of money that Gregory saves over a period of time is a linear model.*
- What are the variables and quantities of this problem?
 - *The rate is \$20 per week, w is the number of weeks, and the initial value is \$500.*
- What function represents the amount of Gregory's money over a period of time (in weeks)?
 - *The model is $f(w) = 20w + 500$. If the function is graphed, the slope of the line is 20 to reflect the constant rate of \$20/week, and the y -intercept is \$500, the initial amount.*

Example 2 (5 minutes)**Example 2**

One of the highlights in a car show event is a car driving up a ramp and flying over approximately five cars placed end-to-end. The ramp is 8 ft. at its highest point, and there is an upward speed of 88 ft/sec before it leaves the top of the ramp. What type of function can best model the height, h , in feet, of the car t seconds after leaving the end of the ramp? Justify your answer mathematically.

- What type of function can best model the height, h , in feet, of the car t seconds after leaving the end of the ramp? What were your clues? Justify your answer mathematically.
 - *Quadratic function. This is an object in motion problem. The car would leave the ramp in an upward and forward motion and then, after travelling higher for a short time, would begin the fall due to the force of gravity.*
- What form would the equation take?
 - *Since the distance is measured in feet and the time in seconds, we would use $h(t) = -16t^2 + v_0t + h_0$, and the equation would be $h(t) = -16t^2 + 88t + 8$.*

Example 3 (5 minutes)**Example 3**

Margie got \$1,000 from her grandmother to start her college fund. She is opening a new savings account and finds out that her bank offers a 2% annual interest rate, compounded monthly. What type of function would best represent the amount of money in Margie's account? Justify your answer mathematically.

- What type of function would best represent the amount of money in Margie's account? Justify your answer mathematically.
 - *Exponential Function. The amount of deposited money grows over time at a constant rate, and the pattern can be best described by an exponential function, $f(x) = ab^x$, where a represents the initial investment, and b is the expression $\left(1 + \frac{r}{n}\right)$ as defined in the compounded interest formula:*

$$P(t) = P_0 \left(1 + \frac{r}{n}\right)^{nt}$$

- What function represents the amount of money deposited in the bank compounded monthly at the rate of 2%, if the initial amount of deposit was \$1,000?

$$A(n) = 1000 \left(1 + \frac{0.02}{12}\right)^{12t}$$

- Note that we do not know how long Margie plans to leave the money in her account, so we do not know what the value of t is yet.

Remind students that percentages, in most cases, must be changed to decimals when used in exponential expressions.

Exercises (12 minutes)

Have students work with a partner or small group to determine the function model for the situation described. Circulate around the room to make sure students understand the exercises. If time is short, some of these might be used for additional homework.

Exercises

- City workers recorded the number of squirrels in a park over a period of time. At the first count, there were 15 pairs of male and female squirrels (30 squirrels total). After 6 months, the city workers recorded a total of 60 squirrels, and after a year, there were 120.

- What type of function can best model the population of squirrels recorded over a period of time, assuming the same growth rate and that no squirrel dies?

Exponential function

- Write a function that represents the population of squirrels recorded over x number of years. Explain how you determined your function.

Students may use the general exponential function $f(x) = ab^x$, by figuring out that this is a doubling exponential problem (in this case, the number of squirrels doubles every 6 months). So, the function would be $f(x) = 30(2)^{2x}$ because the squirrel population would double twice each year.

- A rectangular photograph measuring 8 in by 10 in is surrounded by a frame with a uniform width, x .

- What type of function can best represent the area of the picture and the frame in terms of x (the unknown frame's width)? Explain mathematically how you know.

This function can be best represented by a quadratic function because this is an area problem where the product of two linear measurements results in a quadratic.

- Write an equation in standard form representing the area of the picture and the frame. Explain how you arrive at your equation.

The dimensions of the picture are 8 in by 10 in. Taking into consideration the width of the frame, we have to add $2x$ to both the width and the length of the picture. Doing so results in $(8 + 2x)$ and $(10 + 2x)$. So, the area of the picture and the frame is $A(x) = (8 + 2x)(10 + 2x)$ or $A(x) = 4x^2 + 36x + 80$.

- A ball is tossed up in the air at an initial rate of 50 ft/sec from 5 ft. off the ground.

- What type of function models the height (h , in feet) of the ball after t seconds?

Quadratic function

- b. Explain what is happening to the height of the ball as it travels over a period of time (in t seconds).

The initial height of the ball is 5 ft., and it travels upward with an initial velocity of 50 ft./sec. As time increases, the ball continues to travel upward, with the force of gravity slowing it down, until it reaches the maximum height and falls back to the ground.

- c. What function models the height, h (in feet), of the ball over a period of time (in t seconds)?

$$h(t) = -16t^2 + 50t + 5$$

4. A population of insects is known to triple in size every month. At the beginning of a scientific research project, there were 200 insects.

- a. What type of function models the population of the insects after t years?

Exponential function

- b. Write a function that models the population growth of the insects after t years.

Using the general form for exponential growth ($b = 3$), we have the initial population of 200 and 12 is the number of growth cycles over t years. The function would be $f(t) = 200(3)^{12t}$.

$$P = P_0(1 + r)^{nt}$$

$$P(t) = 200(1 + 2)^{12t}, \text{ where } r \text{ represents the growth rate at } 200\% \text{ and } n = 12.$$

$$\text{So, } P(t) = 200(3)^{12t}.$$

Closing (1 minute)

- How would you know which function to use to model a word problem?
 - *If the word problem talks about repeatedly adding or subtracting a constant value, then it is linear. If the problem involves motion of objects subject to nonzero, constant acceleration (e.g., gravity) over time or represents an area, then it is quadratic. If the problem is about conventional population growth or compounded interest with a constant growth rate, it is exponential.*

Lesson Summary

The following methods can be used to recognize a function type from a word problem:

1. If a problem requires repeated addition or subtraction of a constant value, then it is represented by a linear function.
2. If a problem involves free-falling motion of an object or an area, then it is represented by a quadratic function.
3. If a problem is about population growth or compound interest, then it is represented by an exponential function.

Exit Ticket (2 minutes)

Name _____

Date _____

Lesson 3: Analyzing a Verbal Description

Exit Ticket

Create a model to compare these two texting plans:

- Plan A costs \$15 a month, including 200 free texts. After 200, they cost \$0.15 each.
- Plan B costs \$20 a month, including 250 free texts. After 250, they cost \$0.10 each.

Exit Ticket Sample Solutions

Create a model to compare these two texting plans:

- Plan A costs \$15 a month, including 200 free texts. After 200, they cost \$0.15 each.
- Plan B costs \$20 a month, including 250 free texts. After 250, they cost \$0.10 each.

$$\text{Monthly cost of Plan A: } A(t) = \begin{cases} 15, & \text{if } t \leq 200 \\ 15 + 0.15(t - 200), & \text{if } t > 200 \end{cases} \text{ where } t \text{ represents the number of texts per month}$$

$$\text{Monthly cost of Plan B: } B(t) = \begin{cases} 20, & \text{if } t \leq 250 \\ 20 + 0.10(t - 250), & \text{if } t > 250 \end{cases} \text{ where } t \text{ represents the number of texts per month}$$

Problem Set Sample Solutions

If time allows, the following Problem Set can be used for additional practice. Otherwise, give this Problem Set as homework. Some of these questions go one step further (i.e., they ask an interpretive question), but they do not complete the modeling cycle.

- The costs to purchase school spirit posters are as follows: two posters for \$5, four posters for \$9, six posters for \$13, eight posters for \$17, and so on.

- What type of function would best represent the cost of the total number of posters purchased?

Linear function

- What function represents the cost of the total number of posters purchased? How did you know? Justify your reasoning.

Let x represent the number of school spirit posters. The four ordered pairs indicate a constant rate of change, ($m = 2$), so the equation is $y = 2x + b$. To find b , we need to substitute any ordered pair, say $(2, 5)$: $5 = 2(2) + b$, so $b = 1$. The final equation for the function is $f(x) = 2x + 1$.

- If you have \$40 to spend, write an inequality to find the maximum number of posters you could buy.

$$2x + 1 \leq 40$$

- NYC Sports Gym had 425 members in 2011. Based on statistics, the total number of memberships increases by 2% annually.

- What type of function models the total number of memberships in this situation?

Exponential function

- If the trend continues, what function represents the total number of memberships in n years? How did you know? Justify your reasoning.

$$f(n) = 425(1 + 0.02)^n$$

The initial number of members is 425. The yearly growth rate of 2% means I have to multiply by 1.02 for each year. So, 1.02 will be the common ratio for this exponential function.

3. Derek throws a baseball upward from an initial height of 3 ft . The baseball hits the ground after 2 seconds.

a. What was the initial velocity of the baseball?

$$h(t) = -16t^2 + v_0t + 3$$

$$h(2) = -16(2)^2 + 2v_0 + 3 = 0$$

$$v_0 = 30.5$$

The initial velocity of the baseball was 0.5 ft/sec.

b. What is the function that models the height, h (in feet), of the baseball over a period of time t (in seconds)?

$$h(t) = -16t^2 + 30.5t + 3$$

c. At what time did the baseball reach its maximum height? What was the maximum height of the baseball?

The baseball reached its maximum height after 0.953125 seconds. The maximum height of the baseball was approximately 17.535 feet.